

Extrapyramidal Symptom Scale (EPSS)[©]

Person's Name: _____

WHY MONITOR?: Schizophrenia²⁰ and the use of antipsychotics are independently associated with increased motor effects, which can contribute to non-adherence to medication, and increased morbidity and mortality^{4, 11, 14, 18}.

MINIMUM MONITORING FREQUENCY: For newly initiated medication: baseline, 1, 3 and 12 months. For persons on the same medication > 1 year: q 12 months.

A. MOTOR SIDE EFFECTS

Medical History		<input type="checkbox"/> No relevant motor/neurological history		<input type="checkbox"/> Motor/neurological disorders	
Details:					
Family History in First Degree Relative		<input type="checkbox"/> No relevant motor/neurological history		<input type="checkbox"/> Motor/neurological disorders	
Details:					
Assessment Date (dd/mm/yy)					
Assessment Completed By:					
Current Antipsychotic and Side Effect Medication(s)					
SUBJECTIVE EXPERIENCE (≤ 1 week)	SCORE: 0 = NONE 1 = QUESTIONABLE 2 = MILD 3 = MODERATE 4 = SEVERE				
	Score	Score	Score	Score	Score
Parkinsonism					
Dyskinesia					
Akathisia					
Dystonia					
PARKINSONISM Score right/left sides as indicated	SCORE: 0 = NONE 1 = QUESTIONABLE 2 = MILD 3 = MODERATE 4 = SEVERE				
	Score	Score	Score	Score	Score
Facial expression – reduced					
Hands – tremor (resting)	R	R	R	R	R
	L	L	L	L	L
Hands – tremor (with arms extended, fingers apart)	R	R	R	R	R
	L	L	L	L	L
Hands – bradykinesia	R	R	R	R	R
	L	L	L	L	L
Elbow – rigidity	R	R	R	R	R
	L	L	L	L	L
Gait – abnormality					
DYSKINESIA Score right/left sides as indicated	SCORE: 0 = NONE 1 = QUESTIONABLE 2 = MILD 3 = MODERATE 4 = SEVERE				
	Score	Score	Score	Score	Score
Face and mouth – <i>with activation</i>					
Tongue – <i>with activation</i>					
Upper extremities – arms, hands	R	R	R	R	R
	L	L	L	L	L
Trunk – neck, shoulders, hips					
Lower extremities – ankles/toes	R	R	R	R	R
	L	L	L	L	L
AKATHISIA	SCORE: 0 = NONE 1 = QUESTIONABLE 2 = MILD 3 = MODERATE 4 = SEVERE				
	Score	Score	Score	Score	Score
Observed motor restlessness					
DYSTONIA	SCORE: 0 = NONE 1 = QUESTIONABLE 2 = MILD 3 = MODERATE 4 = SEVERE				
	Score	Score	Score	Score	Score
Observed dystonia					
Details – name affected body part (e.g. head, extremities, trunk):					

B. ISSUE/ACTION/OUTCOME

DATE	ISSUE	ACTION	OUTCOME

RATIONALE: The purpose of the EPSS is to improve the examination and documentation of motor side effects of antipsychotics, allowing for their earlier detection and management.

SCORING PROCEDURE: See anchor point below to guide scoring. Where uncertainty exists about the severity level of a side effect (HJ. questionable v. mild), record the higher of the two scores.

OKDQGVRQDPVSHFWVRIWKHH[DPLQDWLRQVKRØGEHGRQHZLWKSQDVVLYHPRYHPHQWRIWKHSHUVROVUHVSHFWLYHERGSDUW

SUBJECTIVE EXPERIENCE	
Screen	Ask the person: "During the last week, have you..." D noticed any shakes, muscle stiffness, or problems walking? (PARKINSONISM) E noticed any abnormal body movements? (DYSKINESIA) F felt restless or had the need to move even when you didn't want to? (AKATHISIA) G experienced any muscle spasms that lasted at least 1 minute? (DYSTONIA) If yes, explore and record the pertinent details.
PARKINSONISM	Consists of motor disturbances, which include: tremor, impaired gait/posture, postural instability, rigidity, reduced facial expression/speech, and bradykinesia.
Facial expression - reduced	Observe the person's face for reduced facial expression, decreased blinking or parted lips.
Hands – tremor (resting)	Observe the person's hands for a resting tremor, with the person sitting with their elbows resting on their thighs and their hands hanging over their knees.
Hands – tremor (with arms extended, fingers apart)	Observe the person's hands for a tremor, with their palms facing down and arms fully extended with fingers apart.
Hands – bradykinesia	Ask the person to fully open and close their hands, one at a time, in rapid succession, observing for bradykinesia.
Elbow – rigidity	Flex and extend the person's arms, one at a time, with your thumb on their bicep tendon, noting rigidity ("cog-wheel" or "lead pipe").
Gait – abnormality	Observe the person's gait either entering or exiting the room. Note evidence of stooped posture, shuffling gait, decreased arm swing or bradykinesia.
DYSKINESIA	Characterized by movements that are repetitive, purposeless, and involuntary.
Muscles of face and mouth – with activation	While engaging the person in an activation activity (e.g. finger tapping), observe the person's face and mouth, noting any frowning, blinking, grimacing, puckering, repetitive opening and closing of the mouth, clenching of the jaw or lateral movements of the jaw.
Tongue – with activation	While engaging the person in an activation activity (e.g. finger tapping), with the person's mouth open, observe for in and out or lateral movements of the tongue.
Upper extremities – arms, hands	While the person is sitting in a chair, face the patient to observe for evidence of dyskinesias of the: a) Arms and hands. Do NOT include tremor. b) Ankles and/or toes (including inversion/eversion of the foot). c) Neck, shoulders, hips (including rocking, twisting, squirming).
Trunk – neck, shoulders, hips	
Lower extremities – ankles/toes	
AKATHISIA	Consists of subjective feelings of inner restlessness with the urge to move, and/or objective movements such as restless movement of one extremity, changing position, rocking while standing or sitting, lifting feet as if marching on the spot, and inability to sit down for long periods with pacing back and forth.
Observed motor restlessness – lower limbs	The person should be observed (while seated) for a minimum of 5 minutes. A "severe" score should be reserved for persons who are unable to remain seated for the entire 5 minute time period, due to akathisia.
DYSTONIA	Characterized by muscles which are contracted, contorted and often painful, sometimes accompanied by repetitive jerking or twisting movements, resulting in the person's assuming abnormal postures.
Observed dystonia – head, upper and lower extremities, trunk	The person's entire head, neck, limbs and trunk should be observed while sitting or standing. The details of observed dystonias should be recorded.

AUTHORS

Co-created for the Canadian Consortium for Early Intervention in Psychosis by:

Thomas Hastings, MD, FRCPC
 Associate Clinical Professor Department of Psychiatry McMaster University
 Hamilton Ontario
 Lecturer Department of General Psychiatry University of Toronto
 Toronto Ontario
 Address for correspondence: TRPhastingsO@JPDLO.cRP

Phil Tibbo, MD, FRCPC
 Professor and Dr Paul Janssen Chair in Psychotic Disorders Department of Psychiatry
 Dalhousie University
 BOLID| 1RYD 6FRWLD
 Director Nova Scotia Early Psychosis Program
 Halifax, Nova Scotia

5LFKDUG:LOOLDPV0)5&&
 'LUHFWRURI6FKLJRSKUHQLD6HU'VDFRKHU,VODQKQDOWKXVWRULW\
 9LFWRULD%ULWLVKRØELD
 \$MØFVURIHVVRURI3VKRORJ8QLYHUVRV9LFWRULD
 9LFWRULD%ULWLVKRØELD
 ØLQLFDO3URIHVVRU'HSDUWPHQWVWRIØVW8QLYHUVLWRI%ULWLVKRØELD
 9LFWRULD%ULWLVKRØELD



BUNLQVRQLVP

)DFLDO(\$UHVVLRO
ROH QRVRSWR PVSUHVQW
XHVWLRQDEOH XQFOHDLIVRSWRPSUHVHQW
COG PLOGGHFUHDVHLQIDFLDOH\$UHVVLVHQV
BGHUDWH FROVLVWVHWOEWOXQWGDIIHFWEWXVVRPHBVMVZKHJHJPHQWUIUHTXHQW
VWDULQJJDHJUHGXFHGEOLQLOJ
B YHUH QRIDFLDOH\$UHVVLVHQVWVVDULQJJDHJRWDEONHGXFHGEOLQLOJ

WPRU
ROH QRVA PSWRPVSUHVHQW
XHVWLRQDEOH XQFOHDLIVRSWRPSUHVHQW
COG LQIUHTXHQWOSUHVHQWQDGRUVPDOOIQHDPOLWXGHPRYHPHQW
BGHUDWH IUHTXHQWONVHQWZWKPRHJUDWHDPSOLWXGHPRYHPHQWV
B YHUH FROVLQXRXVOSUHVHQWZWKPRGHUDWHWRRODUJHDPOLWXGHPRYHPHQWV

EJLGLWA
ROH QRVA PSWRPVSUHVHQW
XHVWLRQDEOH XQFOHDLIVRSWRPSUHVHQW
COG VRPHUHVLVWDQFHWRSVVLVYHPRYHPHQW
BGHUDWH PRGHUDWHUHVLVWVRSVVLVYHPRYHPHQWEXWHDVLOVPRYHPORPE
6YHUH PDUNGHUHVLVWDQFHWRSVVLVYHPRYHPHQWGLILFXOWWRPRYHPOLPE

DLW
ROH QRVA PSWRPVSUHVHQW
XHVWLRQDEOH XQFOHDLIVRSWRPSUHVHQW
COG PLOGUHGXFHGSOGXODUDUPPRYHPRYHPHQWVWLIISXPHGUMHSOHOJQWK
BGHUDWH PRGHUDWHONHGXFHGSOGXODUDUPPRYHPHQWVWLIISXPHGUMHSOHOJQWK
B YHUH QRSHQG XODUDUPPRYHPHQWPDUNHGOMHGXFHGVVHSHOHOJQWKVKXIILOLQJJDVGLILFXOWW
ZWKWXUQLQJQRWDEOHVWRVWKLBR

NDWKLVLVD

ROH QRVA PSWRPVSUHVHQW
XHVWLRQDEOH XQFOHDLIVRSWRPSUHVHQW
COG UHVWVWPRYHPHQWVQHJHJWUHLVWRPLVWRPLQFUFUHDVHLQSRVLLWRQFQDOJHVEXWDEOHWRUHPDIB
VHDWHGKURXJKGXUDWLRQRIHQFRXQWU
BGHUDWH IUHTXHQWPRYHPHQWRIQHRUPRUHHJWUHLVWYHUNJHTXHQWRSVLLWRQFQDOJHVPD
EHDVNLQJWRJHJXSDQDQONEXWDEOHWRUHPDLOVHDWHGKURXJKGXUDWLRQRIHQFRXQWU
6YHUH QHDFRQVWDQVPRYHPHQWRIPRUHWKQDQRQHJWUHLVWYHUNJHTXHQWRSVLLWRQFQDO
URFNLOJ8DEOHVWRPDLQVHDGRUWKHGXDUDWLRQRIHQFRXQWU

VWRQLD

ROH QRVA PSWRPVSUHVHQW
XHVWLRQDEOH XQFOHDLIVRSWRPSUHVHQW
COG YLVLEOHXPVFOHFRQWUDFWRLOKLFKSDWLHQWVDEOHWRFRUHUHFVWZKPLQLPDOLPSDFWROSRVWXUH
3DWLHQWGRHRVQRWSDSHDUVREHLQDQGLVWUHV
BGHUDWH QRWDEOHQDQGVXVWZKQFOHFRQWUDFWRLOKLFKSDWLHQWVDEOHWRFRUHUHFVWZKPLQLPDOLPSDFWROSRVWXUH
VRUHSHDWHGOWKLPDFWROSRVWXUHBLHQWVDEOHWRFRUHUHFVWZKPLQLPDOLPSDFWROSRVWXUH
B YHUH QRWDEOHQDQGVXVWZKQFOHFRQWUDFWRLOKLFKSDWLHQWVDEOHWRFRUHUHFVWZKPLQLPDOLPSDFWROSRVWXUH
VXEVWDQWLDQVWZKQFOHFRQWUDFWRLOKLFKSDWLHQWVDEOHWRFRUHUHFVWZKPLQLPDOLPSDFWROSRVWXUH

VNLQHLVD

XHVWLRQDEOH XQFOHDLIVRSWRPSUHVHQW
COG LQIUHTXHQWOSUHVHQWPRYHPHQWVWKLKSDWLHQWVDEOHWRFRUHUHFVWZKPLQLPDOLPSDFWROSRVWXUH
RIWLPH
BGHUDWH IUHTXHQWONVHQWZWKPRHJUDWHDPSOLWXGHPRYHPHQWV
B YHUH FROVLQXRXVOSUHVHQWPRYHPHQWVWKLKSDWLHQWVDEOHWRFRUHUHFVWZKPLQLPDOLPSDFWROSRVWXUH
EULHISHULRGVRIWLPH

Adler LA, Angrist B, Reiter S, Rotrosen J. Neuroleptic Induced Akathisia: A Review. J Psychopharmacology. 1989;97:1-11.
Albanese A, Sorbo FD, Comella C, Jinnah HA, Mink JW, Post B, Vidailhet M, Volkman J, Warner TT, Leentjens AFG, et al. Dystonia Rating Scales: Critique and Recommendations. Mov Disord. 2013;28(7):874-883.
Barnes TR. A Rating Scale for Drug-Induced Akathisia. Br. J. Psychiatry. 1989;154:672-676.
Browne S, Roe M, Lane A, Gervin M, Morris M, Kinsella A, Larkin C, Callaghan EO. Quality of Life in Schizophrenia: Relationship to Sociodemographic Factors, Symptomatology and Tardive Dyskinesia. Acta Psychiatr Scand. 1996;94:118-124.
%VOBU KDQGDNDQVROLWRLURUJH[WUDSNDPLGDOGHIIHFVWSDWLHQWV DQWLVSVKRWRUHDWVPHQVSOHWVHGLVFRKH, ULVWRVWZKQDSDVVKRORJLFDOPHGLFLOH 6HS
Chouinard G, Margolese HC. Manual for the Extrapryamidal Symptom Rating Scale (ESRS). Schiz Res. 2005;76:247-265.
Colosimo C, Martinez-Martin P, Fabbrini G, Hauser RA, Merello M, Miyasaki J, Poewe W, Sampaio C, Rascol O, Stebbins GT, et al. Task Force Report on Scales to Assess Dyskinesia in Parkinson's Disease: Critique and Recommendations. Mov Disord. 2010;25(9):1131-1142.
Cortese L, Jog M, McAuley TJ, Kotteda V, Costa G. Assessing and Monitoring Antipsychotic-Induced Movement Disorders in Hospitalized Patients: A Cautionary Study. Can J Psychiatry. 2004;49(1):31-36.
Dean CE, Russell JM, Kuskowski MA, Caligiuri MP, Nugent SM. Clinical Rating Scales and Instruments: How Do They Compare in Assessing Abnormal, Involuntary Movements? J Clin Psychopharmacology. 2004;24(3):298-304.
Fahn S, Elton RL, Members of the UPDRS Development Committee: Unified Parkinson's Disease Rating Scale. In: Fahn S, Marsden CD, Calne DB, Lieberman A, eds: Recent Developments In Parkinson's Disease. Florham Park, NJ: Macmillan Health Care Information 1987;153-163, 293-304.
OHLHacker WW, Meise U, Gunther V, Kurz M. Compliance with Antipsychotic Drug Treatment: Influence of Side Effects. Acta Psychiatr Suppl. 1994;382:11-15.
Gervin M, Barnes TRE. Assessment of Drug-Related Movement Disorders in Schizophrenia. Adv Psychiatr Treat. 2000;6:332-343.
Guy W. ECDEU Assessment Manual for Psychopharmacology. Washington, DC: U.S. Dept. of Health, Education, and Welfare. 1976:534-537.
Hawley CJ, Fineberg N, Roberts AG, Baldwin D, Sahadevan A, Sharman V. The Use of the Simpson Angus Scale for the Assessment of Movement Disorder: A Training Guide. Int J Psychiatry Clin Pract. 2003;7:249-257.
RJDQ 73 ZDG \$ 6MHFWL\$HVSROVWRHROHSWDTQIG2VFRPLQ6FKLJRSKUHQLD \$ 5H(DPLQDWLWRPSDULQJ 7ZROHDVWV 3VKKROHG
.QRO: .HLMVHUV&DOVHGS YDQODUR 5- 6VWHPDWYDQVWRLQD5DWLQFDOHVIRU 'UXQGHG3DUNLQVRQLVPDQGS5HFRPPHQGDWLRQVWVHVDUHFQLO 3VKKRSKDUVDFRQ /RROHQ'R'RUVFKRW& YDQPHUW\$ 2RVWHOERVO806LMEH6 7KH06(6 7HDP7KH 6FKHGQRUWKH\$VHVVPHPDQVQGHG ORYHPHQWVRUGHLASOR' 7HVW5HVVHVW5H0DQDQVH0DQGLVQVWVHRSVKRKRSDUPDFRQ 2 ODUJROHVN KRKODUG* :DQWUHV/DUFRK%HDKRODUJ 5HODWLRQVWVZHHQ 6VRSVKRWRVLF, QGHQWKLVDQGDUGLFWNLQHLVLDQGLDQVFLGDOLQV6FKLJRSKUHQLD ,PSDFVRI QRJDSLOHQDQGDQDQJSDLSORVND3VKKLDVWVHOJ ODWKHZQ*UDWJ 6 SHWZM*HRUJH 9 ODWKHZQ %DVLOS\$WLSVKKRWLF, QGHG ORYHPHQWVRUGHUWDOVWLRQDQGHDPHQW3VKKLDWU\ODUJFK 3DSD 6'DJ]DQ36SROWDQHRX ORYHPHQWLVVWVWLSVKKRWLV1D3DVLHQVWVZVWV)LUVV(SLVRG\$VKKRVHVS 6VWHPDWVWVHLHZ3VKKROOHG 2 6LPSVRQ0 QJX :-6 \$5DWLQJ6FDQHU(WUDSNDPLGDOGHIIHFVWSDWLHQWV 6FDQGP 6WRPVNL10RUULVRO3HNU \$ \$WLSVKKRWVHGLFDWLRQVHGHFVWVHVVPHPHQVRS VVWHPDWVWVHLHZ\$WUDOLDQZ=HDODQG -RQDQR 3VKKLDWU\ ODV

